

Year: 2021

Vol.: 98

Fasc.: 1-2

Title: Recollements associated to cotorsion pairs over upper triangular matrix rings

Author(s): Rongmin Zhu, Yeyang Peng and Nanqing Ding

Let A, B be two rings and $T = \begin{pmatrix} A & M \\ 0 & B \end{pmatrix}$ with M an A - B -bimodule. Suppose that we are given two complete hereditary cotorsion pairs $(\mathcal{A}_A, \mathcal{B}_A)$ and $(\mathcal{C}_B, \mathcal{D}_B)$ in A -Mod and B -Mod, respectively. We define two cotorsion pairs $(\Phi(\mathcal{A}_A, \mathcal{C}_B), \text{Rep}(\mathcal{B}_A, \mathcal{D}_B))$ and $(\text{Rep}(\mathcal{A}_A, \mathcal{C}_B), \Psi(\mathcal{B}_A, \mathcal{D}_B))$ in T -Mod and show that both of these cotorsion pairs are complete and hereditary. If we are given two cofibrantly generated model structures \mathcal{M}_A and \mathcal{M}_B on A -Mod and B -Mod, respectively, then using the result above, we investigate when there exists a cofibrantly generated model structure \mathcal{M}_T on T -Mod and a recollement of $\text{Ho}(\mathcal{M}_T)$ relative to $\text{Ho}(\mathcal{M}_A)$ and $\text{Ho}(\mathcal{M}_B)$. Finally, some applications are given in Gorenstein homological algebra.

Address:

Rongmin Zhu
Department of Mathematics
Nanjing University
Nanjing, 210093
P. R. China

Address:

Yeyang Peng
Department of Mathematics
Nanjing University
Nanjing, 210093
P. R. China

Address:

Nanqing Ding
Department of Mathematics
Nanjing University
Nanjing, 210093
P. R. China